

Relations between *ex falso*, *tertium non datur*, and double negation elimination

Pedro Francisco Valencia Vizcaíno

March 30, 2013

Abstract

We show which implicational relations hold between the three principles *ex falso sequitur quodlibet*, *tertium non datur*, and *double negation elimination*, on the basis of minimal logic.

1 Introduction

By *ex falso sequitur quodlibet*, or simply *ex falso* for short, we mean intuitively the idea that we can derive anything from a contradiction. In slightly more formal terms, we mean a principle that allows us to derive all formulas of the form

$$\neg A \supset (A \supset B)$$

where A and B are arbitrary formulæ.

By *tertium non datur* we mean a principle that allows us to derive all formulas of the form

$$A \vee \neg A$$

where A is an arbitrary formula, and by *double negation elimination* we mean a principle that allows us to derive all formulas of the form

$$\neg\neg A \supset A.$$

The formal system we feel represents minimal logic is the sequent calculus system for intuitionistic logic LJ (see for example [4]) without the rule *weakening:right*. We will argue that this rule really represents the logical principle *ex falso sequitur quodlibet*, and apply a standard cut elimination procedure to this and a slightly modified version of this system to show what relations hold between the logical principles *tertium non datur*, *ex falso*, and *double negation elimination*.

2 Minimal Logic

Let ML stand for the formal system obtained from the formal system for intuitionist logic LJ by removing the rule weakening:right. We are thinking in the context of Sequent Calculus.

Lemma 1 *Over ML it is equivalent to have $\rightarrow \neg A \supset (A \supset B)$ as initial sequents for all formulas A and B and to have the rule weakening:right.*

Proof. If we add weakening:right to ML we can prove ex-falso quodlibet in the following straightforward way:

$$\frac{\frac{\frac{A \rightarrow A}{\neg A, A \rightarrow} \text{w:r}}{\neg A, A \rightarrow B}}{A, \neg A \rightarrow B} \text{w:r} \\ \frac{\neg A \rightarrow A \supset B}{\rightarrow \neg A \supset (A \supset B)}$$

Perhaps a bit less obvious is the fact that with this form of ex-falso quodlibet ML proves *Weakening-right*:

$$\begin{array}{c}
\frac{A \rightarrow A \quad B \rightarrow B}{A \supset B, A \rightarrow B} \\
\frac{\neg A \rightarrow \neg A \quad \frac{A \rightarrow A \quad B \rightarrow B}{A \supset B, A \rightarrow B}}{\neg A \supset (A \supset B), \neg A, A \rightarrow B} \\
\frac{\rightarrow \neg A \supset (A \supset B)}{\neg A, A \rightarrow B} \text{Cut} \\
\frac{A \rightarrow A \quad \frac{A \rightarrow}{\rightarrow \neg A}}{A \rightarrow \neg A} \\
\frac{A \rightarrow A \quad \frac{A \rightarrow \neg A}{A \rightarrow A \wedge \neg A}}{A \rightarrow B} \text{Cut}
\end{array}$$

☐

Lemma 2 *The following hold on the basis of minimal logic:*

1. Double negation elimination implies tertium non datur and ex falso.
2. Tertium non datur + ex falso imply double negation elimination.

Proof.

1. The following is a derivation in minimal logic which shows how to get $A \vee \neg A$ from $\neg\neg(A \vee \neg A) \supset (A \vee \neg A)$:

$$\begin{array}{c}
\frac{A \rightarrow A}{A \rightarrow A \vee \neg A} \\
\frac{\neg(A \vee \neg A), A \rightarrow}{A, \neg(A \vee \neg A) \rightarrow} \\
\frac{\neg(A \vee \neg A) \rightarrow \neg A}{\neg(A \vee \neg A) \rightarrow A \vee \neg A} \\
\frac{\neg(A \vee \neg A), \neg(A \vee \neg A) \rightarrow}{\neg(A \vee \neg A) \rightarrow} \\
\frac{\neg(A \vee \neg A) \rightarrow}{\rightarrow \neg \neg(A \vee \neg A)} \\
\hline
\frac{\rightarrow \neg \neg(A \vee \neg A) \supset (A \vee \neg A) \quad \frac{\neg \neg(A \vee \neg A) \rightarrow \neg \neg(A \vee \neg A) \quad A \vee \neg A \rightarrow A \vee \neg A}{\neg \neg(A \vee \neg A) \supset (A \vee \neg A), \neg \neg(A \vee \neg A) \rightarrow A \vee \neg A}}{\neg \neg(A \vee \neg A) \rightarrow A \vee \neg A} \\
\hline
\rightarrow A \vee \neg A
\end{array}$$

And the next one shows that double negation elimination proves ex falso

$$\begin{array}{c}
\frac{A \rightarrow A}{\neg A, A \rightarrow} \\
\frac{\neg B, \neg A, A \rightarrow}{\neg A, A \rightarrow \neg \neg B} \\
\frac{\neg A, A \rightarrow \neg \neg B \quad B \rightarrow B}{\neg \neg B \supset B, \neg A, A \rightarrow B} \\
\hline
\frac{\rightarrow \neg \neg B \supset B \quad \neg A, A \rightarrow B}{\neg A, A \rightarrow B} \\
\frac{A, \neg A \rightarrow B}{\neg A \rightarrow A \supset B} \\
\hline
\rightarrow \neg A \supset (A \supset B)
\end{array}$$

2. To see that with ex falso and tertium non datur we can obtain double negation elimination we use the following derivation:

$$\begin{array}{c}
\frac{A \rightarrow A}{\neg \neg A, A \rightarrow A} \quad \frac{\neg A \rightarrow \neg A}{\neg \neg A, \neg A \rightarrow} \\
\frac{A, \neg \neg A \rightarrow A}{\neg A, \neg \neg A \rightarrow A} \quad \frac{\neg A, \neg \neg A \rightarrow}{\neg A, \neg \neg A \rightarrow A} \text{ w:r} \\
\hline
\frac{\rightarrow A \vee \neg A \quad A \vee \neg A, \neg \neg A \rightarrow A}{\neg \neg A \rightarrow A}
\end{array}$$

□

2.1 Minimal Logic and Tertium non datur

The way we choose to include *tertium non datur* in ML to get ML^+ is by adding to ML the following inference rule:

$$\frac{A, \Gamma \rightarrow \Delta \quad \neg A, \Gamma \rightarrow \Delta}{\Gamma \rightarrow \Delta}$$

It is easy to see that with this rule we can derive $A \vee \neg A$ for any formula A :

$$\frac{\frac{A \rightarrow A}{A \rightarrow A \vee \neg A} \quad \frac{\neg A \rightarrow \neg A}{\neg A \rightarrow A \vee \neg A}}{\rightarrow A \vee \neg A}$$

It is also easy to see that this rule is perfectly compatible with LK because $A \vee \neg A$ is always derivable in that system:

$$\frac{\frac{\frac{A \rightarrow A}{A \rightarrow A \vee \neg A}}{\rightarrow A \vee \neg A, \neg A} \quad \frac{A, \Gamma \rightarrow \Delta \quad \neg A, \Gamma \rightarrow \Delta}{A \vee \neg A, \Gamma \rightarrow \Delta}}{\Gamma \rightarrow \Delta}$$

So we are somewhat justified in claiming this to be an addition of *tertium non datur* to ML to get ML^+ .

3 Cut Elimination

We will do what Girard does in his book *Proofs and Types* [2] to prove cut elimination for ML and ML^+ :

3.1 Reductions

$$\frac{\frac{A, \Gamma \rightarrow}{\Gamma \rightarrow \neg A} \quad \frac{\Pi \rightarrow A}{\neg A, \Pi \rightarrow}}{\Gamma, \Pi \rightarrow}$$

changes to

$$\frac{\frac{\frac{\Pi \rightarrow A \quad A, \Gamma \rightarrow}{\Gamma, \Pi \rightarrow}}{\Gamma \rightarrow A} \quad \frac{\Gamma \rightarrow B}{\Gamma \rightarrow A \wedge B} \quad \frac{B, \Pi \rightarrow \Lambda}{A \wedge B, \Pi \rightarrow \Lambda}}{\Gamma, \Pi \rightarrow \Lambda}$$

changes to

$$\frac{\frac{\frac{\Gamma \rightarrow B}{\Gamma \rightarrow A \vee B} \quad \frac{A, \Pi \rightarrow \Lambda \quad B, \Pi \rightarrow \Lambda}{A \vee B, \Pi \rightarrow \Lambda}}{\Gamma, \Pi \rightarrow \Lambda}}$$

changes to

$$\frac{\Gamma \rightarrow B \quad B, \Pi \rightarrow \Lambda}{\Gamma, \Pi \rightarrow \Lambda}$$

$$\frac{\frac{A, \Gamma \rightarrow B}{\Gamma \rightarrow A \supset B} \quad \frac{\Pi \rightarrow A \quad B \rightarrow \Lambda}{A \supset B, \Pi \rightarrow \Lambda}}{\Gamma, \Pi \rightarrow \Lambda}$$

changes to

$$\frac{\frac{\Pi \rightarrow A \quad A, \Gamma \rightarrow B}{\Pi, \Gamma \rightarrow B} \quad B \rightarrow \Lambda}{\Pi, \Gamma \rightarrow \Lambda}$$

$$\frac{\frac{\Gamma \rightarrow A(a)}{\Gamma \rightarrow \forall x A(x)} \quad \frac{A(t), \Pi \rightarrow \Lambda}{\forall x A(x), \Pi \rightarrow \Lambda}}{\Gamma, \Pi \rightarrow \Lambda}$$

changes to

$$\frac{\Gamma \rightarrow A(t) \quad A(t), \Pi \rightarrow \Lambda}{\Gamma, \Pi \rightarrow \Lambda}$$

$$\frac{\frac{\Gamma \rightarrow A(t)}{\Gamma \rightarrow \exists x A(x)} \quad \frac{A(a), \Pi \rightarrow \Lambda}{\exists x A(x), \Pi \rightarrow \Lambda}}{\Gamma, \Pi \rightarrow \Lambda}$$

changes to

$$\frac{\Gamma \rightarrow A(t) \quad A(t), \Pi \rightarrow \Lambda}{\Gamma, \Pi \rightarrow \Lambda}$$

The degree $\partial(A)$ of a *formula* is defined as:

- $\partial(A) = 1$ for A atomic
- $\partial(A \wedge B) = \partial(A \vee B) = \partial(A \supset B) = \max(\partial(A), \partial(B)) + 1$
- $\partial(\neg A) = \partial(\forall x A(x)) = \partial(\exists x A(x)) = \partial(A) + 1$

The *degree* of a *cut rule* is defined to be the degree of the formula which it cuts.

The *degree* $d(\pi)$ of a *proof* is the least upper bound (l.u.b.) of the set of degrees of its cut rules. So $d(\pi) = 0$ iff π is cut-free.

The height of a proof is

- $h(\pi) = 1$ if π is an axiom

- $h(\pi) = \text{l.u.b.}(h(\pi_1), h(\pi_2)) + 1$ if π is a proof with a binary last inference rule whose premisses are proved by the subproofs π_1 and π_2 .
- $h(\pi) = h(\pi_0) + 1$ if π is a proof with a unary last inference rule (eg weakening left) and the subproof of the premiss of this inference rule is π_0 .

Notation If Π is a sequence of formulas then by $\Pi - C$ we mean Π where an *arbitrary* number of occurrences of C have been deleted.

Lemma 3 *Let C be a formula of degree n , and π, π' be minimal logic proofs of $\Gamma \rightarrow C$ and $\Pi \rightarrow \Lambda$ of degrees $< n$. Then there is a minimal logic proof of degree $< n$ of $\Gamma, \Pi - C \rightarrow \Lambda$.*

Proof. By induction on $h(\pi) + h(\pi')$.

If both π and π' are proofs of height one, ie, axioms we are in the following situation:

$$C \rightarrow C \quad D \rightarrow D$$

If $D \neq C$ then

$$\frac{D \rightarrow D}{C, D \rightarrow D} \text{w:l}$$

gives what we wanted.

If $D = C$ then either

$$\frac{C \rightarrow C}{C, C \rightarrow C}$$

or

$$C \rightarrow C$$

gives what we wanted.

If π is an axiom and π' isn't then the situation is the following

$$C \rightarrow C \quad \frac{\pi' \vdash}{\Pi \rightarrow \Lambda}$$

and we want to conclude $C, \Pi - C \rightarrow \Lambda$ which we can do by weakening from $\Pi \rightarrow \Lambda$.

If π' is an axiom then the situation is

$$\frac{\pi \vdash}{\Gamma \rightarrow C} \quad D \rightarrow D$$

and again if $D \neq C$ then we get what we want from $D \rightarrow D$ by weakening up to $\Gamma, D \rightarrow D$.

If $D = C$ then $\Gamma \rightarrow C$ or $\Gamma, C \rightarrow C$ give us what we want.

If neither π nor π' are axioms then either both π and π' introduce C in their last inference rule, or one of them doesn't.

If π doesn't, and in particular this includes the rule

$$\frac{A, \Gamma \rightarrow C \quad \neg A, \Gamma \rightarrow C}{\Gamma \rightarrow C}$$

then, we use the induction hypothesis with the premiss(es) of the last inference rule of π and $\Pi \rightarrow \Lambda$, and then use the last inference rule of π . For example:

$$\frac{\frac{A, \Gamma \rightarrow C \quad \Pi \rightarrow \Lambda}{A, \Gamma, \Pi \rightarrow C \rightarrow \Lambda} \quad \frac{\neg A, \Gamma \rightarrow C \quad \Pi \rightarrow \Lambda}{\neg A, \Gamma, \Pi \rightarrow C \rightarrow \Lambda}}{\Gamma, \Pi \rightarrow C \rightarrow \Lambda}$$

Now if both π and π' introduce C in their last inferences then we should look at the reductions above. For example:

If we had

$$\frac{\frac{\Gamma \rightarrow A(t)}{\Gamma \rightarrow \exists x A(x)} \quad \frac{A(a), \Pi \rightarrow \Lambda}{\exists x A(x), \Pi \rightarrow \Lambda}}{\Gamma, \Pi \rightarrow \Lambda}$$

then by induction hypothesis we would have a proof of degree $< n$ of $\Gamma, A(a), \Pi \rightarrow C \rightarrow \Lambda$. Then, by the substitution lemma, we would get (merely by substituting) $\Gamma, A(t), \Pi \rightarrow C \rightarrow \Lambda$. So things look like this:

$$\frac{\Gamma \rightarrow A(t) \quad \frac{\Gamma, A(t), \Pi \rightarrow C \rightarrow \Lambda}{A(t), \Gamma, \Pi \rightarrow C \rightarrow \Lambda}}{\Gamma, \Gamma, \Pi \rightarrow C \rightarrow \Lambda}$$

□

4 Main Results

Lemma 4 *Let B be an atomic formula. If the sequent $\Gamma \rightarrow B$ is provable in ML^+ then there is a formula F which has B as a subformula not occurring in the scope of a negation such that $F \in \Gamma$.*

Proof. By induction on the height of a cut-free proof of $\Gamma \rightarrow B$. If $\Gamma \rightarrow B$ is an axiom then it is the axiom $B \rightarrow B$. In this case $F = B$ works because B is a subformula of F and it does not appear in the scope of a negation (F has no negation symbols at all).

Let π be a proof of $\Gamma \rightarrow B$:

If the last inference rule of π is weakening left

$$\frac{\Gamma' \rightarrow B}{D, \Gamma' \rightarrow B}$$

then by induction hypothesis we have a formula F in Γ' in which B appears as a subformula and not within the scope of a negation. This same formula F appears in the sequence D, Γ' .

If the last inference rule of π was contraction left or exchange left we observe that it is very similar to the first case (weakening left).

If the last rule was our new rule

$$\frac{D, \Gamma \rightarrow B \quad \neg D, \Gamma \rightarrow B}{\Gamma \rightarrow B}$$

then we note that by induction hypothesis, in $\neg D, \Gamma \rightarrow B$ we already have the necessary formula F ; moreover, this formula cannot be $\neg D$ because since B is atomic it cannot be equal to $\neg D$ and it also cannot appear as a subformula of $\neg D$ not inside the scope of a negation, because every proper subformula of $\neg D$ is inside the scope of the head of $\neg D$ which is a negation. This means that the formula F is already in Γ .

The other cases seem easy to verify since we can't use any negation rules if we have the atomic formula B on the right hand side of the sequent, and the other inference rules won't add a negation to any of the formulas in their premisses.

Therefore the statement follows. $\square$

Corollary 1 *If A and B are atomic formulas then*

$$ML^+ \not\vdash \neg A, A \rightarrow B$$

Corollary 2 *ML^+ does not prove weakening:right.*

Corollary 3 *ML^+ does not prove ex falso quodlibet.*

Theorem 1 *The following hold on the basis of minimal logic:*

1. *Double negation elimination implies Ex falso and Tertium non datur.*
2. *Ex falso + Tertium non datur imply Double negation elimination.*
3. *Ex falso does not imply Tertium non datur.*
4. *Tertium non datur does not imply Ex falso.*
5. *Tertium non datur does not imply Double negation elimination.*

References

- [1] Gentzen, Gerhard, *Untersuchungen über das logische Schließen*, Mathematische Zeitschrift. 39 (2): 176-210, 1934.
- [2] Girard, Jean-Yves, *Proofs and Types*, Cambridge University Press, 1989.
- [3] Negri, Sara, von Plato, Jan, *Structural Proof Theory*, Cambridge University Press, 2001.
- [4] Takeuti, Gaisi, *Proof Theory*, North Holland Publishing Company, 1975.
- [5] Troelstra Anne S., Schwichtenberg Helmut, *Basic Proof Theory*, Cambridge University Press, 2000.